

Geometry of the unification of quantum mechanics and relativity*

A. Kryukov

Department of Mathematics, University of Wisconsin Colleges

The paper summarizes and generalizes a recently proposed mathematical framework that unifies the standard formalisms of special relativity and quantum mechanics. The framework is based on Hilbert spaces H of functions of four space-time variables \mathbf{x}, t , furnished with an additional indefinite inner product invariant under Poincaré transformations, and isomorphisms of these spaces that preserve the indefinite metric. The indefinite metric is responsible for breaking the symmetry between space and time variables and for selecting a family of Hilbert subspaces that are preserved under Galileo transformations. Within these subspaces the usual quantum mechanics with Schrödinger evolution and t as the evolution parameter is derived. Simultaneously, the Minkowski space-time is isometrically embedded into H , Poincaré transformations have unique extensions to isomorphisms of H and the embedding commutes with Poincaré transformations. The main new result is a proof that the framework accommodates arbitrary pseudo-Riemannian space-times furnished with the action of the diffeomorphism group.

I. PRELIMINARIES

Resolving the tension between quantum theory and relativity is one of the most important problems of modern physics. In this paper the issue is presented mathematically and the goal is to find a formalism that unifies the mathematics used in both theories. More specifically, the challenge is to find a meaningful way of “encoding” differential geometry of finite dimensional pseudo-Riemannian manifolds into the theory of infinite dimensional Hilbert spaces. Provided such a “unified” formalism exists, it may be useful in particular, in studying the issues of compatibility of quantum theory and relativity and the problem of emergence of the classical world in quantum theory.

The first thing that comes to mind is that the theory of representations of groups provides a partial answer to the challenge. Indeed, it allows one to represent symmetries of a physical system in terms of linear transformations on the Hilbert space of states of the system. For instance, symmetries of Minkowski space-time can be represented by unitary transformations on the Hilbert space of states of a relativistic system. However, despite the undeniable significance of representations in physics there is more to the problem than representations of groups can provide. For instance, if the symmetry group (i.e., group of isometries) of a Riemannian manifold is trivial, representations of the group do not contain any information about the manifold.

On the other hand, there is a celebrated Gel’fand-Kolmogorov theorem Ref.[1] that states that an arbitrary compact

* This paper is based on a talk given at IARD 2008 conference. An expanded version of a part of this paper can be found in Ref.[7]

Hausdorff space X is homeomorphic to the space of all evaluation homomorphisms (delta functions) in the infinite-dimensional vector space dual to the Banach algebra $C(X)$ of continuous functions on X . In other words, X can be identified with the set of all delta functions in the space dual to $C(X)$. This theorem provides a link between the topology of space X and algebra $C(X)$ and may be useful to tackle the problem. However, $C(X)$ is not a Hilbert space, the topology is poorer than the Riemannian structure and the condition of compactness is too restrictive. In addition, the fact that elements of $C(X)$ are functions on X makes it difficult to use $C(X)$ independently of X . This is a problem if one has in mind the goal of deriving the classical from the quantum theory.

In the case of a single particle system in \mathbb{R}^3 the most obvious physically meaningful embedding of \mathbb{R}^3 into the space of states of a particle is by identifying a point \mathbf{a} in \mathbb{R}^3 with the state $\delta_{\mathbf{a}}^3(\mathbf{x}) = \delta^3(\mathbf{x} - \mathbf{a})$ of the particle found at \mathbf{a} . This embedding was usefully explored in Refs.[2]-[6] to develop a geometric approach to quantum mechanics. Results of Ref.[7] prove that an embedding of this kind is also ideally suited for addressing the issues of the unification of quantum mechanics and special relativity. In the present paper the results of Ref.[7] are summarized, updated and extended to include arbitrary space-time manifolds.

II. QUANTUM MECHANICS AND SPECIAL RELATIVITY

The idea of embedding \mathbb{R}^3 into the Hilbert space of states of a particle via the correspondence between points $\mathbf{a} \in \mathbb{R}^3$ and states $\delta_{\mathbf{a}}^3$ runs into an immediate difficulty. Namely, delta functions are not in the common space $L_2(\mathbb{R}^3)$ of Lebesgue square-integrable functions on \mathbb{R}^3 . So to use this correspondence one first needs to find a Hilbert space of functions that contains delta functions and that is “approximately equal” to the space $L_2(\mathbb{R}^3)$. The following theorem takes care of this task (see Ref.[7] for proofs of theorems in this section).

Theorem 1. *The Hilbert space \mathbf{H} obtained by completing the space $L_2(\mathbb{R}^3)$ in the metric defined by the inner product*

$$(\varphi, \psi)_{\mathbf{H}} = \left(\frac{L}{\sqrt{2\pi}} \right)^3 \int e^{-\frac{L^2}{2}(\mathbf{x}-\mathbf{y})^2} \varphi(\mathbf{x}) \bar{\psi}(\mathbf{y}) d^3\mathbf{x} d^3\mathbf{y} \quad (1)$$

with a positive constant L contains delta functions and their derivatives. Furthermore, for a sufficiently large L the \mathbf{H} and L_2 -norms of any given function $f \in L_2(\mathbb{R}^3)$ are arbitrarily close to each other.

The map $\omega : a \rightarrow \delta_{\mathbf{a}}^3$ is one-to-one, so the set \mathbb{R}^3 can be identified with the set of all delta functions in \mathbf{H} . But is it possible to induce the manifold structure and the Euclidean metric on the image $M_3 = \omega(\mathbb{R}^3) \subset \mathbf{H}$ from the metric on \mathbf{H} ? The following theorem gives a positive answer.

Theorem 2. *The map $\omega : a \rightarrow \delta_{\mathbf{a}}^3$ is an isometric embedding of the space \mathbb{R}^3 with the Euclidean metric into the space \mathbf{H} defined in theorem 1.*

Note that the map ω is not linear. In particular, because the norm of any delta function $\delta_{\mathbf{a}}^3$ in \mathbf{H} is the same, the image $\omega(\mathbb{R}^3)$ is a submanifold of the sphere in \mathbf{H} . So the vector structure on \mathbb{R}^3 is not compatible with the vector

structure on \mathbf{H} . However, one can introduce a vector structure on the image M_3 by defining the operations of addition \oplus and multiplication by a scalar $\lambda \odot$ via $\omega(a) \oplus \omega(b) = \omega(a + b)$ and $\lambda \odot \omega(a) = \omega(\lambda a)$. Moreover, because ω is a homeomorphism onto M_3 , these operations are continuous in the topology of $M_3 \subset \mathbf{H}$.

The Euclidean space \mathbb{R}^3 is now a submanifold in the Hilbert space \mathbf{H} , formed by point-supported state functions $\delta_{\mathbf{a}}^3$, and with the Euclidean metric induced by the embedding. In this sense the Riemannian structure of the Euclidean space is encoded into \mathbf{H} . At the same time, the space \mathbf{H} is approximately equal to the space $L_2(\mathbb{R}^3)$ (symbolically, $\mathbf{H} \approx L_2(\mathbb{R}^3)$). That is, provided the constant L in theorem 1 is sufficiently large, the \mathbf{H} -norms of typical square-integrable functions will be as close as we wish to their $L_2(\mathbb{R}^3)$ norms. Accordingly, if \mathbf{H} is used in place of $L_2(\mathbb{R}^3)$ in the usual quantum mechanics, the expected values, probabilities of transition and other measured quantities remain practically the same, ensuring consistency with experiment.

The next step is to extend results of theorems 1, 2 to Minkowski space-time and build a framework that incorporates relativity on Minkowski space-time and non-relativistic quantum mechanics of a single particle. To obtain a relativistic framework one needs to work with spaces of functions of four variables \mathbf{x}, t . The following theorem generalizes the results of theorem 1 to this case.

Theorem 3. *Let \tilde{H} be the Hilbert space of functions f of four variables $x = (\mathbf{x}, t)$ that is the completion of the space $L_2(\mathbb{R}^4)$ in the metric given by the kernel $e^{-\frac{1}{2}(x-y)^2}$. Let \tilde{H}_{ev} be the subspace of all functions in \tilde{H} that are even in the time variable t . Likewise, let \tilde{H}_{odd} be the subspace of all functions that are odd in t . Then \tilde{H} is the orthogonal sum of \tilde{H}_{ev} and \tilde{H}_{odd} . Furthermore, consider the set H of all functions $f(\mathbf{x}, t) = e^{-t^2} \varphi(\mathbf{x}, t)$ with $\varphi \in \tilde{H}$. Consider the Hermitian form $(f, g)_{H_\eta}$ on H given by*

$$(f, g)_{H_\eta} = \int e^{-\frac{1}{2}(\mathbf{x}-\mathbf{y})^2 + \frac{1}{2}(t-s)^2} f(\mathbf{x}, t) \bar{g}(\mathbf{y}, s) d^3 \mathbf{x} dt d^3 \mathbf{y} ds \quad (2)$$

and let $(f, f)_{H_\eta} \equiv \|f\|_{H_\eta}^2$ be the corresponding quadratic form, or the squared H_η -norm. Then H is exactly the set of functions whose even and odd components have a finite H_η -norm. Moreover, H furnished with the inner product $(f, g)_{H_+} = (\varphi, \psi)_{\tilde{H}}$, where $f(\mathbf{x}, t) = e^{-t^2} \varphi(\mathbf{x}, t)$, $g(\mathbf{x}, t) = e^{-t^2} \psi(\mathbf{x}, t)$ is a Hilbert space. The Hermitian form (2) defines an indefinite, non-degenerate inner product on H , such that $\|f\|_{H_\eta}^2 > 0$ for all even functions $f \neq 0$ and $\|f\|_{H_\eta}^2 < 0$ for all odd functions $f \neq 0$ in H . Finally, H contains the delta functions $\delta_a^4(x) = \delta^4(x - a)$ and their derivatives.

The space H is an example of what is called the *Krein space*. A Krein space is a complex vector space V with a Hermitian inner product $(f, g)_V$ and such that V is the direct sum of two spaces H_1, H_2 that are Hilbert with respect to the inner products $(f, g)_V$ and $-(f, g)_V$ respectively and that are orthogonal in the inner product on V . The reason for the particular Krein space in the theorem to be useful becomes clear from the following analogue of theorem 2.

Theorem 4. *The map $\omega : N \rightarrow H$, $\omega(a) = \delta_a^4$ is an embedding that identifies the Minkowski space N with the submanifold M_4 of H of all delta functions δ_a^4 , $a \in N$. Under the embedding the indefinite metric on H yields the Minkowski metric on M_4 , while the \tilde{H} -metric yields the ordinary Euclidean metric on M_4 .*

So, similarly to the space \mathbb{R}^3 , the Minkowski space N is now encoded into the space H . As before, the map ω is not linear, but the image M_4 can be furnished with a linear structure, induced from N .

With N embedded into H , the essential question now is how to “lift” the theory of relativity from N onto H . The following simple result turns out to be the key to resolving the issue and explaining the geometry of the embedding of N into H .

Theorem 5. *The set M_4 is a complete set in the Hilbert spaces H and \tilde{H} . Accordingly, linear combinations of elements of M_4 span these spaces. The elements of any finite subset of M_4 are linearly independent. No two elements of M_4 are orthogonal in the indefinite metric or the Hilbert metric on H , or the \tilde{H} -metric. However, the inner product $(\delta_a^4, \delta_b^4)_{\tilde{H}}$ with $a = (\mathbf{a}, a_0)$ and $b = (\mathbf{b}, b_0)$ tends to zero when the Euclidean norm $\|a - b\|_{\mathbb{R}^4}$ tends to infinity. Similarly, the indefinite inner product $(\delta_a^4, \delta_b^4)_{H_\eta}$ tends to zero when the Minkowski norm $\|a - b\|_N^2 = (\mathbf{a} - \mathbf{b})^2 - (a_0 - b_0)^2$ tends to infinity.*

Before formulating the next result, let’s make a few observations. If Π is a Poincaré transformation, and f is a function(al) in H , then $\delta_\Pi : f \longrightarrow f \circ \Pi^{-1}$ is a linear map on H . Because the Hilbert metric on H is not invariant under general Poincaré transformations, the operator δ_Π is not bounded as a map into H so that the map $\Pi \longrightarrow \delta_\Pi$ is not a representation of the Poincaré group P . However, the set of all functions $f \circ \Pi^{-1}$ with a fixed Π and $f \in H$ form a Hilbert space H' with the inner product defined by $(\delta_\Pi f, \delta_\Pi g)_{H'_+} = (f, g)_{H_+}$. The map $\delta_\Pi : H \longrightarrow H'$ is then an isomorphism of Hilbert spaces. Hilbert spaces obtained in such a way can be thought of as different realizations of one and the same abstract Hilbert space \mathbf{S} . A particular isomorphism $\Gamma : \mathbf{S} \longrightarrow H$ can be thought of as a coordinate chart on \mathbf{S} Ref.[2]. With this in mind one can formulate the following essential result. In the theorem the expression *isometric embedding* refers to an embedding that preserves the indefinite metric. Likewise *isomorphism* is an isomorphism of Hilbert spaces that in addition preserves the indefinite metric.

Theorem 6. *Let $\Gamma : \mathbf{S} \longrightarrow H$ be an isomorphism of the abstract Hilbert space \mathbf{S} with an additional indefinite metric onto the space H of functions defined in theorem 3. Let $\gamma : N \longrightarrow \mathbb{R}^{1,3}$ be a global coordinate chart from the Minkowski space-time onto the coordinate space of the observer in an inertial reference frame K . Let Π be a Poincaré transformation that relates coordinates associated with frames K and K' . Then there exists a unique isometric embedding Ω and a unique isomorphism $\delta_\Pi : H \longrightarrow H'$ such that the diagram*

$$\begin{array}{ccccc}
 \mathbf{S} & \xrightarrow{\Gamma} & H & \xrightarrow{\delta_\Pi} & H' \\
 \uparrow \Omega & & \uparrow \omega & & \uparrow \omega \\
 N & \xrightarrow{\gamma} & \mathbb{R}^{1,3} & \xrightarrow{\Pi} & \mathbb{R}^{1,3}
 \end{array} \tag{3}$$

is commutative. It follows that within the assumptions of the theorem the embedding ω preserves the structure of special relativity and extends it in a unique way to the abstract Hilbert space \mathbf{S} .

A couple of remarks.

1. Note that δ_{Π} maps delta functions to delta functions, so that in accord with the diagram (3) the image of the manifold M_4 in H is the submanifold M_4 in H' . This together with the fact that ω is an isometric embedding is what allows for the usual theory of relativity on Minkowski space-time to be a part of the new framework. At the same time completeness of the set M_4 together with linear independence of its elements (theorem 5) makes the entire construction rigid, ensuring uniqueness of the extension.
2. The proposed method of extension of the Poincaré group action from N onto \mathbf{S} can be applied to *any* group acting on N . Notice that an arbitrary non-linear transformation acting continuously on N becomes linear when extended to \mathbf{S} . That is so because moving across N corresponds to going “across dimensions” of H (theorem 5) so that a linear extension of the transformation becomes possible. Completeness of the set M_4 in H ensures then that such an extension is unique.
3. The Poincaré transformations Π and their extensions $\delta\Pi$ in the theorem are “passive” transformations i.e., they describe changes in coordinate realizations of the fixed, invariant spaces N and \mathbf{S} . The corresponding “active” version of the theorem is also possible and is given by the following analogue of diagram (3):

$$\begin{array}{ccccc}
 \mathbf{S}' & \xleftarrow{\delta_{\Pi}} & \mathbf{S} & \xrightarrow{\Gamma} & H \\
 \uparrow \Omega & & \uparrow \Omega & & \uparrow \omega \\
 N & \xleftarrow{\Pi} & N & \xrightarrow{\gamma} & \mathbb{R}^{1,3}
 \end{array} \tag{4}$$

Here the maps γ, Γ, ω and the embedding $\Omega : N \rightarrow \mathbf{S}$ are the same as before. The Poincaré transformation Π maps Minkowski space N onto itself. The space \mathbf{S}' contains the subset $\Omega(N)$ as a complete set and is otherwise defined by the diagram. The isomorphism δ_{Π} is the linear extension of the map $\Omega\Pi\Omega^{-1}$ from $\Omega(N)$ onto \mathbf{S} .

Having constructed an extension of special relativity to the Hilbert space \mathbf{S} let's now turn to the embedding of quantum mechanics into the same framework. The first thing to do is to relate the Hilbert space H of functions of four variables \mathbf{x}, t to the usual Hilbert spaces of functions of three variables \mathbf{x} with t as a parameter of evolution. For this consider the family of subspaces H_{τ} of H each consisting of all functionals $\varphi_{\tau}(\mathbf{x}, t) = \psi(\mathbf{x}, t)\delta(t - \tau)$ for some fixed $\tau \in \mathbb{R}$.

Theorem 7. *Under the inclusion $i : H_{\tau} \rightarrow H$ the indefinite inner product on H yields a Hilbert metric on H_{τ} for all $\tau \in \mathbb{R}$. The inner products H_{+} and \tilde{H} on H yield an equivalent Hilbert metric on H_{τ} . Furthermore, let $\mathbf{H} \approx L_2(\mathbb{R}^3)$ be the Hilbert space defined in theorem 1 (with $L = 1$ and a sufficiently small scale to make the approximation valid). Then for all $\tau \in \mathbb{R}$ the map $I : H_{\tau} \rightarrow \mathbf{H}$ defined by $I(\varphi_{\tau})(\mathbf{x}) = \psi(\mathbf{x}, \tau)$ is an isomorphism of Hilbert spaces.*

The map I basically identifies each subspace H_{τ} with the usual space $L_2(\mathbb{R}^3)$ of state functions on \mathbb{R}^3 considered at time τ . The reason why these particular subspaces are physically meaningful becomes clear from the following result that relates the dynamics on the family of subspaces H_{τ} and the usual space $L_2(\mathbb{R}^3)$ of states of a spinless non-relativistic particle.

Theorem 8. Let $\hat{h} = D + V(\mathbf{x}, t)$ be a Hamiltonian, such that D is a differential operator in the spatial coordinates and V is a function. Then the path $\varphi_\tau(\mathbf{x}, t) = \psi(\mathbf{x}, t)\delta(t - \tau)$ in H satisfies the equation $\frac{d\varphi_\tau}{d\tau} = \left(-\frac{\partial}{\partial t} - i\hat{h}\right)\varphi_\tau$ if and only if the function $\psi(\mathbf{x}, t)$ satisfies the Schrödinger equation $\frac{\partial\psi(\mathbf{x}, t)}{\partial t} = -i\hat{h}\psi(\mathbf{x}, t)$. At each point of the path φ_τ the components $-i\hat{h}\varphi_\tau$, $-\frac{\partial\varphi_\tau}{\partial t}$ of the velocity vector $\frac{d\varphi_\tau}{d\tau}$ are orthogonal in the indefinite inner product, \tilde{H} -inner product and the inner product on the space H_T of the time co-moving representation.

Here the space H_T of the time co-moving representation is defined by application of the isomorphism

$$(\delta_{\Pi_\tau} f)(\mathbf{x}, t) = f(\mathbf{x}, t - \tau) \quad (5)$$

to the space H (i.e., the representation is the map $\delta_{\Pi_\tau} \circ \Gamma$, where Γ is the same as in theorem 6). The theorem claims that the ordinary Schrödinger evolution can be recovered from the evolution φ_τ in the space H of functions of four variables by projecting the path φ_τ onto the “co-moving” subspace H_τ identified via I with $\mathbf{H} \approx L_2(\mathbb{R}^3)$. While the component $-i\hat{h}\varphi_\tau$ of the velocity describes the motion within the subspace H_τ , the orthogonal (“vertical”) component $-\frac{\partial\varphi_\tau}{\partial t}$ of the velocity is due to the motion of the subspace H_τ itself.

Several important observations must be made at this point.

1. Under integration in time the time variable gets replaced with the parameter τ . In other words, for motions within the family H_τ the evolution parameter τ used to describe motions in the space H of functions of four variables becomes identified with the usual time variable that appears in Schrödinger equation.
2. The delta factor $\delta(t - \tau)$ in functions in H_τ removes integration in time and therefore eliminates the effect of interference in time that is present for more general elements of H . In fact, the norm of superposition $\psi_1(\mathbf{x}, t)\delta(t - \tau) + \psi_2(\mathbf{x}, t)\delta(t - \tau)$ of functions in H_τ in either H_η , \tilde{H} , or H_T -metrics is equal to $\|\psi_1(\mathbf{x}, \tau) + \psi_2(\mathbf{x}, \tau)\|_{\mathbf{H}}$, which approximates the standard expression due to the relationship $\mathbf{H} \approx L_2(\mathbb{R}^3)$.
3. The space H was needed to identify Minkowski space with an isometrically embedded submanifold $M_4 \subset H$. If this embedding is accepted, the reason for the delta factor $\delta(t - \tau)$ in the non-relativistic limit has a simple explanation. In fact, elements of the space H have the form $e^{-t^2}\varphi(\mathbf{x}, t)$, where φ is in the space $\tilde{H} \approx L_2(\mathbb{R}^4)$ (and the meaning of approximation is the same as in theorem 1). Likewise, the space H_T of the time co-moving representation defined by Eq.(5) consists of the functions $e^{-(t-\tau)^2}\varphi(\mathbf{x}, t)$, with $\varphi \in \tilde{H} \approx L_2(\mathbb{R}^4)$. The variables \mathbf{x}, t enter symmetrically in the definition of $L_2(\mathbb{R}^4)$, while the factor $e^{-(t-\tau)^2}$ breaks the symmetry between \mathbf{x} and t by making a typical element of H_T well localized in the time variable. In a sufficiently small scale the factor $e^{-(t-\tau)^2}$ as a function of $t - \tau$ quickly falls off to almost zero and can be replaced with the delta function $\delta(t - \tau)$. This yields the set of functions in the family of spaces H_τ and by theorems 7 and 8 allows for the usual formalism of quantum mechanics.
4. Subspaces H_τ are not preserved under the maps δ_{Π} in theorem 6. In fact, δ_{Π} mixes space and time coordinates and therefore does not preserve the form $\varphi(\mathbf{x}, t)\delta(t - \tau)$ of elements of H_τ in general. This is not surprising

because standard quantum mechanics is non-relativistic. However, to provide a valid foundation of the non-relativistic quantum mechanics these subspaces must be preserved under Galileo transformations. A Galileo transformation G yields the map $\delta_G : H \rightarrow H'$ defined by $\delta_G f = f \circ G^{-1}$ for all $f \in H$. This map transforms the state $\varphi(\mathbf{x}, t)\delta(t - \tau)$ into the state $\varphi(A\mathbf{x} + \mathbf{v}t + \mathbf{b}, t + c)\delta(t + c - \tau)$, where A is an orthogonal transformation, \mathbf{v}, \mathbf{b} are 3-vectors and c is a real number. Recall now that φ is an element of the Hilbert space \mathbf{H} with metric given by the kernel $e^{-\frac{1}{2}(\mathbf{x}-\mathbf{y})^2}$. This kernel is obviously invariant under Galileo transformations so that the function $\varphi(A\mathbf{x} + \mathbf{v}t + \mathbf{b}, t + c)$ is still an element of \mathbf{H} . One concludes that Galileo transformations yield isomorphisms between subspaces H_τ (and that the map $G \rightarrow \delta_G$, where δ_G is considered as acting on \mathbf{H} is a unitary representation of the Galileo group).

5. The equation $\frac{d\varphi_\tau}{d\tau} = \left(-\frac{\partial}{\partial t} - i\hat{h}\right)\varphi_\tau$ with usual Hamiltonian is a well known non-relativistic limit of the Stueckelberg-Schrödinger equation in the theory of Stueckelberg Ref.[8] and Horwitz & Piron Ref.[9]. This theory treats space and time symmetrically and predicts interference in time Refs.[10],[11]. The non-relativistic limit of Stueckelberg theory was investigated by Horwitz and Rotbart Ref.[12]. The approximate equality of the time variable t with the evolution parameter τ obtained in Ref.[12] is consistent with the definition of H_τ .
6. Newton and Wigner Ref.[13] argue that delta functions δ_a^4 cannot represent spatially localized states in a relativistic theory. However, their derivation is based on the condition of orthogonality of a localized state and its spatial displacement, which is not valid in the proposed framework. Newton and Wigner mention other difficulties with delta states: (1) they are not square integrable, (2) if a sequence of square integrable functions converges to δ_a^4 in one frame, it will not do so in general in a Lorentz transformed frame. However, the delta functions do have a finite norm in the new framework. Also, if a sequence of functions approaches δ_a^4 in the space H , then, according to the diagram (3), the Lorentz transformed sequence will approach $\delta_{\Gamma a}^4$ in the space H' of the frame K' . The operator having δ_a^4 as an eigenfunction with the eigenvalue a^μ is the operator \widehat{x}^μ of multiplication by the variable x^μ , $\mu = 0, 1, 2, 3$. Note that the delta function locality is present in the relativistic Stueckelberg theory, which is off-shell. If the Stueckelberg expectation value of the dynamical variable \widehat{x}^μ is decomposed into a direct integral over mass, then for each definite mass in the integral, the Newton-Wigner operator (having Newton-Wigner localized states as eigenstates) emerges. Locality is restored in the result of the integral Refs.[9],[12]. The covariant property of the states δ_a^4 and the operator \widehat{x}^μ does not mean by itself that the found objects are physical. There are well known difficulties: (1) the wave packet $\delta_{\mathbf{a}}^3$ contains negative energy components; (2) if such a packet is allowed to evolve by the usual relativistic equations it will evolve out of the light cone Ref.[14]. Although these difficulties are typical for relativistic on-shell wave equations and were understood within the Stueckelberg approach Ref.[9], they must be reexamined in the new setting.

The following theorem completes the embedding of the standard quantum mechanical formalism into the framework.

Theorem 9. *An arbitrary isomorphism $V : H \rightarrow H'$ of Hilbert spaces that preserves the indefinite metric on H*

and maps subspaces H_τ of functions $\psi(\mathbf{x}, t)\delta(t - \tau)$ onto themselves yields a family of unitary transformations on the space $\mathbf{H} \approx L_2(\mathbb{R}^3)$ of functions $\psi(\mathbf{x}, t)$. Moreover, all unitary transformations on \mathbf{H} can be obtained in such a way.

III. GENERALIZING THE FRAMEWORK TO CURVED SPACE-TIME MANIFOLDS

So far the discussion involved only the classical 3-dimensional Euclidean space and the Minkowski space-time. If the approach is taken seriously, it becomes essential to check its validity for more general space-times N . It is also important to see whether the Hilbert space into which N is embedded can be defined without specifying the manifold first. For manifolds without additional (pseudo-) Riemannian structure the issues are resolved by the following theorem.

Theorem 10. *Given an arbitrary real n -dimensional manifold N there exists a Hilbert space $H_{\mathbb{R}^n}$ of continuous functions on \mathbb{R}^n , such that the set M_n of all delta functions in the dual space $H_{\mathbb{R}^n}^*$ is an embedded submanifold of $H_{\mathbb{R}^n}^*$ diffeomorphic to N .*

In essence, the theorem claims that an arbitrary n -dimensional manifold can be “encoded” into an appropriate Hilbert space of functions on \mathbb{R}^n . To get an idea of how to find the Hilbert space $H_{\mathbb{R}^n}$, especially when the topology of the manifold is not trivial, consider the case of a circle S^1 . In this case the space $H_{\mathbb{R}}$ must be a Hilbert space of continuous functions on \mathbb{R} . To ensure that the image M_1 of the map $\omega : \mathbb{R} \rightarrow H_{\mathbb{R}}^*$, $\omega(a) = \delta_a$ is a circle, one needs $\delta_a = \delta_{a+2\pi}$ for all $a \in \mathbb{R}$, which means that functions in $H_{\mathbb{R}}$ must be 2π -periodic. To satisfy these conditions, consider the Sobolev space of continuous 2π -periodic functions on \mathbb{R} with the inner product $(f, g) = \int_{-\pi}^{\pi} (f(x)\bar{g}(x) + f'(x)\bar{g}'(x)) dx$. It is easy to check that the set of all delta functions in the dual space $H_{\mathbb{R}}^*$ with the induced topology is homeomorphic to the circle S^1 .

A particular manifold in the theorem is encoded by fixing the *functional content* of the Hilbert space rather than fixing the domain of the functions. To put it differently, the manifold M_n is “made of” functions and not points in the domain of the functions.

The problem of isometric embeddings of Riemannian and pseudo-Riemannian manifolds is now handled by the following theorem.

Theorem 11. *Let N be a Riemannian or pseudo-Riemannian smooth manifold of dimension n . For any point $x \in N$ there is a neighborhood W of x in N and a Hilbert or Krein space H that contains delta functions (evaluation functionals) $\delta_a^{(n)}$ for all a in an open set U in \mathbb{R}^n such that the set M_n of all these delta functions is an embedded submanifold of H isometric to W .*

Proof. It is known that an arbitrary smooth Riemannian or pseudo-Riemannian manifold N of dimension n admits an isometric embedding into the Euclidean or pseudo-Euclidean space \mathbb{R}^p of a sufficiently large dimension $p \geq n$, $p = k + l$, where (k, l) is the signature of the metric on \mathbb{R}^p . Also, by an obvious generalization of theorems 3 and 4

the map $\Omega : \mathbb{R}^p \longrightarrow H_p$, $\Omega(A) = \delta_A^{(p)}$ is an isometric embedding of the space \mathbb{R}^p into the Krein space H_p defined via the inner product

$$(f, g)_{H_p} = \int e^{-\frac{1}{2}((X^1-Y^1)^2+\dots+(X^k-Y^k)^2)+\frac{1}{2}((X^{k+1}-Y^{k+1})^2+\dots+(X^p-Y^p)^2)} f(X^1, \dots, X^p) \bar{g}(Y^1, \dots, Y^p) d^p X d^p Y \quad (6)$$

with $d^p X = dX^1 \dots dX^p$, $d^p Y = dY^1 \dots dY^p$. Note that the analogues of H_{ev} , H_{odd} in theorem 3 are obtained here by representing an arbitrary function $f(X^1, \dots, X^k, X^{k+1}, \dots, X^p)$ as the sum of “even”

$$\frac{1}{2} (f(X^1, \dots, X^k, X^{k+1}, \dots, X^p) + f(X^1, \dots, X^k, -X^{k+1}, \dots, -X^p)) \quad (7)$$

and “odd”

$$\frac{1}{2} (f(X^1, \dots, X^k, X^{k+1}, \dots, X^p) - f(X^1, \dots, X^k, -X^{k+1}, \dots, -X^p)) \quad (8)$$

components. Otherwise the proof mimics the one given in Ref.[7].

Let's form a Hilbert (Krein) subspace H_n of H_p in the following fashion. Let $x \in N$ be a point and let $X^q(u)$, $q = 1, \dots, p$, $u \in U$, and $U \subset \mathbb{R}^n$ be functions describing the isometric embedding of a neighborhood $W \subset N$ of x into \mathbb{R}^p . By permuting indices of the coordinates X^1, \dots, X^p and considering a smaller neighborhood W if necessary one can always ensure that u^1, \dots, u^n are just the first n of the coordinates. So, consider the set H_n of all function(al)s in H_p that have the form $\varphi(X^1, \dots, X^p) \delta(X^{n+1} - X^{n+1}(u)) \dots \delta(X^p - X^p(u))$, or more briefly $\varphi(X) \delta^{(p-n)}(X - X(u))$ with $u \in U$. Denoting the kernel of the metric in H_p , given by Eq.(6), by $k(X, Y)$, we have for the inner product of two such functionals:

$$\int k(X, Y) \varphi(X) \delta^{(p-n)}(X - X(u)) \bar{\psi}(Y) \delta^{(p-n)}(Y - Y(v)) d^p X d^p Y, \quad (9)$$

where $Y^1 = v^1, \dots, Y^n = v^n$. The delta functions remove integration with respect to X^{n+1}, \dots, X^p and Y^{n+1}, \dots, Y^p , which gives

$$\int k(X(u), Y(v)) \varphi(X(u)) \bar{\psi}(Y(v)) d^n u d^n v, \quad (10)$$

where $du = du^1 \dots du^n = dX^1 \dots dX^n$ and similarly for dv . The set H_n is a closed subspace in H_p so it is a Hilbert space. Expression (10) shows that H_n is isomorphic to the Hilbert space H of all functions $\chi(u) = \varphi(X(u))$, $u \in U$ for which $\varphi(X) \delta^{(p-n)}(X - X(u))$ is in H_p , furnished with the inner product $(\chi, \rho)_H = \int k(X(u), Y(v)) \chi(u) \rho(v) d^n u d^n v$.

Obviously, the functionals $\varphi(u) = \delta^{(n)}(u - a)$, $a \in U$ are in H . It remains to show that the metric induced on the set M_n of all such functionals in H is the given (pseudo) Riemannian metric on N . For this consider a curve $u^\mu = a^\mu(\tau)$ in U and the corresponding curve $\varphi_\tau(u) = \delta^{(n)}(u - a(\tau))$ in M_n . For the squared H -norm of the velocity vector $d\delta^{(n)}(u - a(\tau))/d\tau$ we have

$$\int k(X(u), Y(v)) \frac{d\delta^{(n)}(u - a(\tau))}{d\tau} \frac{d\delta^{(n)}(v - a(\tau))}{d\tau} d^n u d^n v. \quad (11)$$

Simplifying this by the chain rule

$$\frac{d\delta^{(n)}(u - a(\tau))}{d\tau} = - \frac{\partial \delta^{(n)}(u - a(\tau))}{\partial u^\mu} \frac{da^\mu(\tau)}{d\tau} \quad (12)$$

followed by integration by parts (see Ref.[2] for justification), one obtains the expression

$$\left. \frac{\partial^2 k(X(u), Y(v))}{\partial u^\mu \partial v^\nu} \right|_{u=v=a(\tau)} = \frac{da^\mu(\tau)}{d\tau} \frac{da^\nu(\tau)}{d\tau}. \quad (13)$$

But

$$\frac{\partial^2 k(X(u), Y(v))}{\partial u^\mu \partial v^\nu} = \frac{\partial^2 k(X(u), Y(v))}{\partial X^r \partial Y^s} \frac{\partial X^r}{\partial u^\mu} \frac{\partial Y^s}{\partial v^\nu}, \quad (14)$$

and for the kernel $k(X, Y)$ given by Eq.(6) one also has

$$\left. \frac{\partial^2 k(X(u), Y(v))}{\partial X^r \partial Y^s} \right|_{u=v} = \eta_{rs}, \quad (15)$$

where η_{rs} are components of the indefinite (Minkowski-like) metric of signature (k, l) on \mathbb{R}^p . So the squared norm of the velocity vector in Eq.(11) is equal to

$$g_{\mu\nu} \frac{da^\mu}{d\tau} \frac{da^\nu}{d\tau}, \quad (16)$$

where

$$g_{\mu\nu} = \eta_{rs} \left. \frac{\partial X^r}{\partial u^\mu} \frac{\partial Y^s}{\partial v^\nu} \right|_{u=v} \quad (17)$$

are the components of the induced metric on M_n .

Recall now that the functions $X^r(u)$ describe the isometric embedding of $W \subset N$ into \mathbb{R}^p . In other words, components of the (pseudo-) Riemannian metric on W are given by

$$\tilde{g}_{\mu\nu} = \eta_{rs} \frac{\partial X^r}{\partial u^\mu} \frac{\partial X^s}{\partial u^\nu}. \quad (18)$$

Since this expression coincides with Eq.(17), the obtained embedding of W into H is isometric. This completes the proof. \square

Several useful observations must be made.

1. Theorem 11 makes it possible to extend the results of theorem 4 to neighborhoods in arbitrary pseudo-Riemannian space-times. In this case the Poincaré group acting on Minkowski space-time is replaced by the group of diffeomorphisms of a particular neighborhood. This yields the following analogue of diagram (3)

$$\begin{array}{ccccc} \mathbf{S} & \xrightarrow{\Gamma} & H & \xrightarrow{\delta_D} & H' \\ \uparrow \Omega & & \uparrow \omega & & \uparrow \omega \\ W & \xrightarrow{\gamma} & U & \xrightarrow{D} & U \end{array} \quad (19)$$

Here W is a neighborhood in curved space-time as defined in theorem 11, γ is a chart on W and U is the corresponding set in \mathbb{R}^4 , D is an arbitrary diffeomorphism of U and δ_D is its extension to the space H constructed in theorem 11. As already mentioned in the remarks following theorem 6, the existence and uniqueness of the isomorphism δ_D and the space H' can be proved as before.

2. Recall that the set M_4 is invariant under transformations δ_{Π} , making it possible to “separate” special relativity from the Hilbert space framework. In the discussion that followed theorem 8 it was verified that the “Galileo maps” δ_G map subspaces H_{τ} onto themselves. This explains why the non-relativistic quantum mechanics could also be developed within a single Hilbert space of functions of three variables. Diagrams (3), (4) provide us with a “covariant” extension of special relativity. Likewise, diagram (19) together with its active version yield a local geometric extension of general relativity. Those extensions are based on isomorphisms of separable Hilbert spaces. If such a scheme is adopted in physics, that would mean that specific functional realizations of the abstract Hilbert space \mathbf{S} , at least within the considered class of realizations, are not physical but rather are similar to various choices of coordinates on space-time. One may disregard this point by saying that considered isomorphisms of Hilbert spaces of functions are direct analogues of well known changes in representation in quantum theory. However, unlike changes of representation that are simply passive changes in the description of physical reality, the transformations considered here can be realized actively. Active transformations create a new physical reality and make the construction physically meaningful.
3. If the discussed embedding of the classical space \mathbb{R}^3 into \mathbf{H} as well as the embeddings of Minkowski space-time and local embeddings of arbitrary curved space-times into the corresponding Hilbert spaces are taken seriously, then the linearity of quantum theory appears in a completely new light. In fact, the geometry of the abstract Hilbert space \mathbf{S} and its realizations like H is linear. It is the non-linearity of submanifolds M_3 and M_4 that seems to be responsible for the non-linear way in which classical world appears to us. By replacing the restricted, “space-time based” view of the world with its extension to the space \mathbf{S} one can perhaps obtain a tool for reconciliation of quantum theory and relativity.

IV. ACKNOWLEDGMENTS

I would like to thank Malcolm Forster for numerous valuable discussions and useful suggestions. I am indebted to Larry Horwitz for critical review of the results, useful references and for a fruitful discussion of the issues of relativistic quantum mechanics. I am grateful to participants of the conference IARD 2008, where this paper was presented for their interest, comments and questions. This work was supported by the NSF grant SES-0849418.

-
- [1] I.M. Gel’fand and A.N. Kolmogorov, *Dokl. Akad. Nauk SSSR*, **22** 11 (1939)
- [2] A. Kryukov, *Int. J. Math. & Math. Sci.* **14**, 2241 (2005)
- [3] A. Kryukov, *Found. Phys.* **34**, 1225 (2004)
- [4] A. Kryukov, *Found. Phys.* **36**, 175 (2006)
- [5] A. Kryukov, *Found. Phys.* **37**, 3 (2007)
- [6] A. Kryukov, *Phys. Lett. A* **370**, 419 (2007)

- [7] A. Kryukov, *J. Math. Phys.* **49**, 102108 (2008)
- [8] E.C.G. Stueckelberg, *Helv. Phys. Acta* **14**, 372, 585 (1941); *Helv. Phys. Acta* **15**, 23 (1942)
- [9] L.P. Horwitz and C. Piron, *Helv. Phys. Acta* **46**, 316 (1973)
- [10] L.P. Horwitz and Y. Rabin, *Lett. Nuovo Cimento* **17**, 501 (1976)
- [11] L.P. Horwitz, *Phys. Lett. A* **355**, 1 (2006)
- [12] L.P. Horwitz and F. Rotbart, *Phys. Rev. D* **24**, 2127 (1981)
- [13] T.D. Newton and E.P. Wigner, *Reviews of Modern Physics*, **21** 400 (1949)
- [14] G.C. Hegerfeldt, *Phys. Rev. D* **10**, 3320 (1974)